arXiv:1403.1601v2 [math.CO] 1 Feb 2016

A bound on the number of edges in graphs without an even cycle

Boris Bukh* Zilin Jiang†,*

Abstract

We show that, for each fixed k, an n-vertex graph not containing a cycle of length $2k$ has at most $80\sqrt{k\log k}\cdot n^{1+1/k}+O(n)$ edges.

MSC CLASSES: 05C35, 05D99, 05C38

Introduction

Let $\mathrm{ex}(n,F)$ be the largest number of edges in an n-vertex graph that contains no copy of a fixed graph F. The systematic study of $\mathrm{ex}(n,F)$ was started by Turán [17] over 70 years ago, and it has developed into a central problem in extremal graph theory (see surveys [9, 10, 15]).

The function $\mathrm{ex}(n,F)$ exhibits a dichotomy: if F is not bipartite, then $\mathrm{ex}(n,F)$ grows quadratically in n, and is fairly well understood. If F is bipartite, $\mathrm{ex}(n,F)$ is subquadratic, and for very few F the order of magnitude is known. The simplest classes of bipartite graphs are trees, complete bipartite graphs, and cycles of even length. Most of the study of $\mathrm{ex}(n,F)$ for bipartite F has been concentrated on these two classes. In this paper, we address the even cycles. For an overview of the status of $\mathrm{ex}(n,F)$ for complete bipartite graphs see [2]. For a thorough survey on bipartite Turán problems see [9].

The first bound on the problem is due to Erdős [5] who showed that $\mathrm{ex}(n,C_4)=O(n^{3/2})$. Thanks to the works of Erdős and Rényi [7], Brown [4, Section 3], and Kövari, Sós and Turán [11] it is now known that

$$\mathrm{ex}(n,C_4)=(1/2+o(1))n^{3/2}.$$

The current best bounds for $\mathrm{ex}(n,C_6)$ for large values of n are

$$0.5338n^{4/3}<\mathrm{ex}(n,C_6)\le 0.6272n^{4/3}$$

due to Füredi, Naor and Verstraëte [8].

A general bound of $\mathrm{ex}(n,C_{2k})\le\gamma_k n^{1+1/k}$, for some unspecified constant γ_k, was asserted by Erdős [6, p. 33]. The first proof was by Bondy and Simonovits [3, Lemma 2], who showed that $\mathrm{ex}(n,C_{2k})\le 20kn^{1+1/k}$ for all sufficiently large n. This was improved by Verstraëte [18] to $8(k-1)n^{1+1/k}$ and by Pikhurko [14] to $(k-1)n^{1+1/k}+O(n)$. The principal result of the present paper is an improvement of these bounds.

*Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh, PA 15213. Supported in part by U.S. taxpayers through NSF grant DMS-1301548.

†Supported in part by U.S. taxpayers through NSF grant DMS-1201380.

Main Theorem. *Suppose G is an n-vertex graph that contains no C_{2k}, and $n \geq (2k)^{8k^2}$ then*

$$\mathrm{ex}(n, C_{2k}) \leq 80\sqrt{k \log k} \cdot n^{1+1/k} + 10k^2 n.$$

It is our duty to point out that the improvement offered by the Main Theorem is of uncertain value because we still do not know if $\Theta(n^{1+1/k})$ is the correct order of magnitude for $\mathrm{ex}(n, C_{2k})$. Only for $k = 2, 3, 5$ are constructions of C_{2k}-free graphs with $\Omega(n^{1+1/k})$ edges known [1, 19, 12, 13]. The first author believes it to be likely that $\mathrm{ex}(n, C_{2k}) = o(n^{1+1/k})$ for all large k. We stress again that the situation is completely different for odd cycles, where the value of $\mathrm{ex}(n, C_{2k+1})$ is known exactly for all large n [16].

Proof method and organization of the paper Our proof is inspired by that of Pikhurko [14]. Apart from a couple of lemmas that we quote from [14], the present paper is self-contained. However, we advise the reader to at least skim [14] to see the main idea in a simpler setting.

Pikhurko's proof builds a breadth-first search tree, and then argues that a pair of adjacent levels of the tree cannot contain a Θ-graph[1]. It is then deduced that each level must be at least $\delta/(k-1)$ times larger than the previous, where δ is the (minimum) degree. The bound on $\mathrm{ex}(n, C_{2k})$ then follows. The estimate of $\delta/(k-1)$ is sharp when one restricts one's attention to a pair of levels.

In our proof, we use three adjacent levels. We find a Θ-graph satisfying an extra technical condition that permits an extension of Pikhurko's argument. Annoyingly, this extension requires a bound on the *maximum degree*. To achieve such a bound we use a modification of breadth-first search that avoids the high-degree vertices.

What we really prove in this paper is the following.

Theorem 1. *Suppose $k \geq 4$, and suppose G is a biparite n-vertex graph of minimum degree at least $2d + 5k^2$, where*

$$d \geq \max(20\sqrt{k \log k} \cdot n^{1/k}, (2k)^{8k}), \tag{1}$$

then G contains C_{2k}.

The Main Theorem follows from Theorem 1 and two well-known facts: every graph contains a bipartite subgraph with half of the edges, and every graph of average degree d_{avg} contains a subgraph of minimum degree at least $d_{\mathrm{avg}}/2$.

The rest of the paper is organized as follows. We present our modification of breadth-first search in Section 1. In Section 2, which is the heart of the paper, we explain how to find Θ-graphs in triples of consecutive levels. Finally, in Section 3 we assemble the pieces of the proof.

1 Graph exploration

Our aim is to have vertices of degree at most Δd for some $k \ll \Delta \ll d^{1/k}$. The particular choice is fairly flexible; we choose to use

$$\Delta \stackrel{\mathrm{def}}{=} k^3.$$

[1] We recall the definition of a Θ-graph in Section 2

Let G be a graph, and let x be any vertex of G. We start our exploration with the set $V_0 = \{x\}$, and mark the vertex x as explored. Suppose $V_0, V_1, \ldots, V_{i-1}$ are the sets explored in the 0th, 1st,...,$(i-1)$st steps respectively. We then define V_i as follows:

1. Let V_i' consist of those neighbors of V_{i-1} that have not yet been explored. Let Bg_i be the set of those vertices in V_i' that have more than Δd unexplored neighbors, and let $\mathrm{Sm}_i = V_i' \setminus \mathrm{Bg}_i$.

2. Define
$$V_i = \begin{cases} V_i' & \text{if } |\mathrm{Bg}_i| > \frac{1}{2k}|V_i'|, \\ \mathrm{Sm}_i & \text{if } |\mathrm{Bg}_i| \leq \frac{1}{2k}|V_i'|. \end{cases}$$
The vertices of V_i are then marked as explored.

We call sets $V_0, V_1, \ldots$ *levels* of G. A level V_i is *big* if $|\mathrm{Bg}_i| > \frac{1}{2k}|V_i'|$, and is *normal* otherwise.

Lemma 2. *If $\delta \leq \Delta d$, and G is a bipartite graph of minimum degree at least δ, then each $v \in V_{i+1}$ has at least δ neighbors in $V_i \cup V_{i+2}'$.*

Proof. Fix a vertex $v \in V(G)$. We will show, by induction on i, that if $v \notin V_1 \cup \cdots \cup V_i$, then v has at least δ neighbors in $V(G) \setminus (V_1 \cup \cdots \cup V_{i-1})$. The base case $i = 1$ is clear. Suppose $i > 1$. If $v \in \mathrm{Bg}_i$, then v has $\Delta d \geq \delta$ neighbors in the required set. Otherwise, v is not in V_i' and hence has no neighbors in V_{i-1}. Hence, v has as many neighbors in $V(G) \setminus (V_1 \cup \cdots \cup V_{i-1})$ as in $V(G) \setminus (V_1 \cup \cdots \cup V_{i-2})$, and our claim follows from the induction hypothesis.

If $v \in V_{i+1}$, then the neighbors of v are a subset of $V_1 \cup \cdots \cup V_i \cup V_{i+2}'$. Hence, at least δ of these neighbors lie in $V_i \cup V_{i+2}'$. □

Trilayered graphs We abstract out the properties of a triple of consecutive levels into the following definition. A *trilayered graph* with layers V_1, V_2, V_3 is a graph G on a vertex set $V_1 \cup V_2 \cup V_3$ such that the only edges in G are between V_1 and V_2, and between V_2 and V_3. If $V_1' \subset V_1$, $V_2' \subset V_2$ and $V_3' \subset V_3$, then we denote by $G[V_1', V_2', V_3']$ the trilayered subgraph induced by three sets V_1', V_2', V_3'. Because the graph G that has been explored is bipartite, there are no edges inside each level. Therefore any three sets V_{i-1}, V_i, V_{i+1}' from the exploration process naturally form a trilayered graph; these graphs and their subgraphs are the only trilayered graphs that appear in this paper.

We say that a trilayered graph has *minimum degree* at least $[A : B, C : D]$ if each vertex in V_1 has at least A neighbors in V_2, each vertex in V_2 has at least B neighbors in V_1, each vertex in V_2 has at least C neighbors in V_3, and each vertex in V_3 has at least D neighbors in V_2. A schematic drawing of such a graph is on the right.



2 Θ-graphs

A Θ-*graph* is a cycle of length at least $2k$ with a chord. We shall use several lemmas from the previous works.

Lemma 3 (Lemma 2.1 in [14], also Lemma 2 in [18]). *Let F be a Θ-graph and $1 \leq l \leq |V(F)| - 1$. Let $V(F) = W \cup Z$ be an arbitrary partition of its vertex set into two non-empty parts such that every path in F of length l that begins in W necessarily ends in W. Then F is bipartite with parts W and Z.*

Lemma 4 (Lemma 2.2 in [14]). *Let $k \geq 3$. Any bipartite graph H of minimum degree at least k contains a Θ-graph.*

Corollary 5. *Let $k \geq 3$. Any bipartite graph H of average degree at least $2k$ contains a Θ-graph.*

For a graph G and a set $Y \subset V(G)$ let $G[Y]$ denote the graph induced on Y. For disjoint $Y, Z \subset V(G)$ let $G[Y, Z]$ denote the bipartite subgraph of G that is induced by the bipartition $Y \cup Z$.

Well-placed Θ-graphs Suppose G is a trilayered graph with layers V_1, V_2, V_3. We say that a Θ-graph $F \subset G$ is *well-placed* if each vertex of $V(F) \cap V_2$ is adjacent to some vertex in $V_1 \setminus V(F)$. The condition ensures that, for each vertex v of F in V_2 there exists a path from the root to v that avoids F.

Lemma 6. *Suppose G is a trilayered graph with layers V_1, V_2, V_3 such that the degree of every vertex in V_2 is at least $2d + 5k^2$, and no vertex in V_2 has more than Δd neighbors in V_3. Suppose t is a nonnegative integer, and let $F = \frac{d \cdot e(V_1, V_2)}{8k|V_3|}$. Assume that*

$$\begin{aligned} &a) && F \geq 2, \\ &b) && e(V_1, V_2) \geq 2kF|V_1|, \\ &c) && e(V_1, V_2) \geq 8k(t+1)^2(2\Delta k)^{2k-1}|V_1|, \\ &d) && e(V_1, V_2) \geq 8(et/F)^t k|V_2|, \\ &e) && e(V_1, V_2) \geq 20(t+1)^2|V_2|. \end{aligned} \tag{2}$$

Then at least one of the following holds:

I) There is a Θ-graph in $G[V_1, V_2]$.

II) There is a well-placed Θ-graph in $G[V_1, V_2, V_3]$.

The proof of Lemma 6 is in two parts: finding trilayered subgraph of large minimum degree (Lemmas 7 and 8), and finding a well-placed Θ-graph inside that trilayered graph (Lemma 9).

Finding a trilayered subgraph of large minimum degree The disjoint union of two bipartite graphs shows that a trilayered graph with many edges need not contain a trilayered subgraph of large minimum degree. We show that, in contrast, if a trilayered graph contains no Θ-graph between two of its levels, then it must contain a subgraph of large minimum degree. The next lemma demonstrates a weaker version of this claim: it leaves open a possibility that the graph contains a denser trilayered subgraph. In that case, we can iterate inside that subgraph, which is done in Lemma 8.

Lemma 7. *Let a, A, B, C, D be positive real numbers. Suppose G is a trilayered graph with layers V_1, V_2, V_3 and the degree of every vertex in V_2 is at least $d + 4k^2 + C$. Assume also that*

$$a \cdot e(V_1, V_2) \geq (A + k + 1)|V_1| + B|V_2|. \tag{3}$$

Then one of the following holds:

I) There is a Θ-graph in $G[V_1, V_2]$.

II) There exist non-empty subsets $V_1' \subset V_1$, $V_2' \subset V_2$, $V_3' \subset V_3$ such that the induced trilayered subgraph $G[V_1', V_2', V_3']$ has minimum degree at least $[A : B, C : D]$.

III) There is a subset $\widetilde{V}_2 \subset V_2$ such that $e(V_1, \widetilde{V}_2) \geq (1 - a)e(V_1, V_2)$, and $|\widetilde{V}_2| \leq D|V_3|/d$.

Proof. We suppose that alternative (I) does not hold. Then, by Corollary 5, the average degree of every subgraph of $G[V_1, V_2]$ is at most $2k$.

Consider the process that aims to construct a subgraph satisfying (II). The process starts with $V_1' = V_1$, $V_2' = V_2$ and $V_3' = V_3$, and at each step removes one of the vertices that violate the minimum degree condition on $G[V_1', V_2', V_3']$. The process stops when either no vertices are left, or the minimum degree of $G[V_1', V_2', V_3']$ is at least $[A : B, C : D]$. Since in the latter case we are done, we assume that this process eventually removes every vertex of G.

Let R be the vertices of V_2 that were removed because at the time of removal they had fewer than C neighbors in V_3'. Put

$$E' \stackrel{\text{def}}{=} \{uv \in E(G) : u \in V_2,\ v \in V_3,\ \text{and } v \text{ was removed before } u\},$$
$$S \stackrel{\text{def}}{=} \{v \in V_2 : v \text{ has at least } 4k^2 \text{ neighbors in } V_1\}.$$

Note that $|E'| \leq D|V_3|$. We cannot have $|S| \geq |V_1|/k$, for otherwise the average degree of the bipartite graph $G[V_1, S]$ would be at least $\frac{4k}{1+1/k} \geq 2k$. So $|S| \leq |V_1|/k$.

The average degree condition on $G[V_1, S]$ implies that

$$e(V_1, S) \leq k(|V_1| + |S|) \leq (k + 1)|V_1|.$$

Let u be any vertex in $R \setminus S$. Since it is connected to at least $(d + 4k^2 + C) - 4k^2 = d + C$ vertices of V_3, it must be adjacent to at least d edges of E'. Thus,

$$|R \setminus S| \leq |E'|/d \leq D|V_3|/d.$$

Assume that the conclusion (III) does not hold with $\widetilde{V}_2 = R \setminus S$. Then $e(V_1, R \setminus S) < (1 - a)e(V_1, V_2)$. Since the total number of edges between V_1 and V_2 that were removed due to the minimal degree conditions on V_1 and V_2 is at most $A|V_1|$ and $B|V_2|$ respectively, we conclude that

$$\begin{aligned} e(V_1, V_2) &\leq e(V_1, S) + e(V_1, R \setminus S) + A|V_1| + B|V_2| \\ &< (k + 1)|V_1| + (1 - a)e(V_1, V_2) + A|V_1| + B|V_2|, \end{aligned}$$

implying that

$$a \cdot e(V_1, V_2) < (A + k + 1)|V_1| + B|V_2|.$$

The contradiction with (3) completes the proof. ∎

Remark. The next lemma can be eliminated at the cost of obtaining the bound $\mathrm{ex}(n, C_{2k}) = O(k^{2/3}n^{1+1/k})$ in place of $\mathrm{ex}(n, C_{2k}) = O(\sqrt{k \log k} \cdot n^{1+1/k})$. To do that, we can set $B \approx k^{2/3}$, $D \approx k^{1/3}$ and $a = 1/2$. One can then show that when applied to trilayered graphs arising from the exploration process the alternative (III) leads to a subgraph of average degree $2k$. The two remaining alternatives are dealt by Corollary 5 and Lemma 9. However, it is possible to obtain a better bound by iterating the preceding lemma.

Lemma 8. *Let C be a positive real number. Suppose G is a trilayered graph with layers V_1, V_2, V_3, and the degree of every vertex in V_2 is at least $d + 4k^2 + C$. Let $F = \frac{d \cdot e(V_1,V_2)}{8k|V_3|}$, and assume that F and $e(V_1, V_2)$ satisfy (2) for some integer $t \geq 1$. Then one of the following holds:*

I) There is a Θ-graph in $G[V_1, V_2]$.

II) There exist numbers A, B, D and non-empty subsets $V_1' \subset V_1$, $V_2' \subset V_2$, $V_3' \subset V_3$ such that the induced trilayered subgraph $G[V_1', V_2', V_3']$ has minimum degree at least $[A : B, C : D]$, with the following inequalities that bind A, B, and D:

$$\begin{aligned} &B \geq 5, \qquad (B-4)D \geq 2k, \\ &A \geq 2k(\Delta D)^{D-1}. \end{aligned} \tag{4}$$

Proof. Assume, for the sake of contradiction, that neither (I) nor (II) hold. With hindsight, set $a_j = \frac{1}{t-j+1}$ for $j = 0, \ldots, t-1$. We shall define a sequence of sets $V_2 = V_2^{(0)} \supseteq V_2^{(1)} \supseteq \cdots \supseteq V_2^{(t)}$ inductively. We denote by

$$d_i \stackrel{\text{def}}{=} e(V_1, V_2^{(i)})/|V_2^{(i)}|$$

the average degree from $V_2^{(i)}$ into V_1. The sequence $V_2^{(0)}, V_2^{(1)}, \ldots, V_2^{(t)}$ will be constructed so as to satisfy

$$e(V_1, V_2^{(i+1)}) \geq (1 - a_i)e(V_1, V_2^{(i)}), \tag{5}$$

$$d_{i+1} \geq d_i \cdot F a_i \prod_{j=0}^{i}(1 - a_j). \tag{6}$$

Note that (5) and the choice of $a_0, \ldots, a_i$ imply that

$$e(V_1, V_2^{(i)}) \geq \tfrac{1}{t+1} e(V_1, V_2). \tag{7}$$

The sequence starts with $V_2^{(0)} = V_2$. Assume $V_2^{(i)}$ has been defined. We proceed to define $V_2^{(i+1)}$. Put

$$\begin{aligned} A &= a_i e(V_1, V_2^{(i)})/2|V_1| - k - 1, \\ B &= a_i d_i/4 + 5, \\ D &= \min(2k, 8k/a_i d_i). \end{aligned}$$

With help of (7) and (2c) it is easy to check that the inequalities (4) hold for this choice of constants.

In addition,

$$\begin{aligned}(A+k+1)|V_1| + B|V_2^{(i)}| &= \tfrac{3}{4}a_i e(V_1, V_2^{(i)}) + 5|V_2^{(i)}| \\ &\overset{(2e)}{\leq} \tfrac{3}{4}a_i e(V_1, V_2^{(i)}) + \tfrac{1}{4(t+1)^2}e(V_1, V_2) \\ &\overset{(7)}{\leq} a_i e(V_1, V_2^{(i)}).\end{aligned}$$

So, the condition (3) of Lemma 7 is satisfied for the graph $G[V_1, V_2^{(i)}, V_3]$. By Lemma 7 there is a subset $V_2^{(i+1)} \subset V_2^{(i)}$ satisfying (5) and

$$|V_2^{(i+1)}| \leq D|V_3|/d.$$

Next we show that the set $V_2^{(i+1)}$ satisfies inequality (6). Indeed, we have

$$\begin{aligned}d_{i+1} = \frac{e(V_1, V_2^{(i+1)})}{|V_2^{(i+1)}|} &\geq \frac{(1-a_i)e(V_1, V_2^{(i)})}{D|V_3|/d} \geq (1-a_i)a_i d_i \frac{d}{8k|V_3|}e(V_1, V_2^{(i)}) \\ &\overset{(5)}{\geq} (1-a_i)a_i d_i \frac{d \cdot e(V_1, V_2)}{8k|V_3|}\prod_{j=0}^{i-1}(1-a_j) = d_i \cdot F a_i \prod_{j=0}^{i}(1-a_j).\end{aligned}$$

Iterative application of (6) implies

$$d_t \geq d_0 F^t \prod_{j=0}^{t-1} a_j(1-a_j)^{t-j} \geq d_0 F^t \prod_{j=0}^{t-1}\frac{e^{-1}}{t-j+1} = d_0\frac{(F/e)^t}{(t+1)!}. \tag{8}$$

If we have $|V_2^{(t)}| < |V_1|$, then the average degree of induced subgraph $G[V_1, V_2^{(t)}]$ is greater than $e(V_1, V_2^{(t)})/|V_1| \overset{(7)}{\geq} e(V_1, V_2)/(t+1)|V_1| \overset{(2c)}{\geq} 2k$, which by Corollary 5 leads to outcome (I).

If $|V_2^{(t)}| \geq |V_1|$ and $d_t \geq 4k$, then the average degree of $G[V_1, V_2^{(t)}]$ is at least $d_t/2 \geq 2k$ because d_t is the average degree of $V_2^{(t)}$ into V_1, again leading to the outcome (I). So, we may assume that $d_t < 4k$. Since $(t+1)! \leq 2t^t$ we deduce from (8) that

$$d_0 < 4k(t+1)!(e/F)^t \leq 8k(et/F)^t.$$

This contradicts (2d), and so the proof is complete. □

Locating well-placed Θ-graphs in trilayered graphs We come to the central argument of the paper. It shows how to embed well-placed Θ-graphs into trilayered graphs of large minimum degree. Or rather, it shows how to embed well-placed Θ-graphs into regular trilayered graphs; the contortions of the previous two lemmas, and the factor of $\sqrt{\log k}$ in the final bound, come from authors' inability to deal with irregular graphs.

Lemma 9. *Let A, B, D be positive real numbers. Let G be a trilayered graph with layers V_1, V_2, V_3 of minimum degree at least $[A : B, d + k : D]$. Suppose that no vertex in V_2 has more than Δd neighbors in V_3. Assume also that*

$$B \geq 5 \tag{9}$$

$$(B-4)D \geq 2k-2 \tag{10}$$

$$A \geq 2k(\Delta D)^{D-1}. \tag{11}$$

Then G contains a well-placed Θ-graph.

Proof. Assume, for the sake of contradiction, that G contains no well-placed Θ-graphs. Leaning on this assumption we shall build an arbitrary long path P of the form



where, for each i, vertices v_i and v_{i+1} are joined by a path of length $2D$ that alternates between V_2 and V_3. Since the graph is finite, this would be a contradiction.

While building the path we maintain the following property:

$$\text{Every } v \in P \cap V_2 \text{ has at least one neighbor in } V_1 \setminus P. \tag{$\star$}$$

We call a path satisfying ($\star$) *good*.

We construct the path inductively. We begin by picking v_0 arbitrarily from V_1. Suppose a good path $P = v_0 \rightsquigarrow v_1 \rightsquigarrow \cdots \rightsquigarrow v_{l-1}$ has been constructed, and we wish to find a path extension $v_0 \rightsquigarrow v_1 \rightsquigarrow \cdots \rightsquigarrow v_{l-1} \rightsquigarrow v_l$.

There are at least about A ways to extend the path by a single vertex. The idea of the following argument shows that many of these extensions can be extended to another vertex, and then another, and so on.

For each $i = 1, 2, \ldots, 2D-1$ we shall define a family $\mathcal{Q}_i$ of good paths that satisfy

1. Each path in $\mathcal{Q}_i$ is of the form $v_0 \rightsquigarrow v_1 \rightsquigarrow \cdots \rightsquigarrow v_{l-1} \rightsquigarrow u$, where $v_{l-1} \rightsquigarrow u$ is a path of length i that alternates between V_2 and V_3. The vertex u is called a *terminal* of the path. The set of terminals of the paths in $\mathcal{Q}_i$ is denoted by $T(\mathcal{Q}_i)$. Note that $T(\mathcal{Q}_i) \subset V_2$ for odd i and $T(\mathcal{Q}_i) \subset V_3$ for even i.

2. For each i, the paths in $\mathcal{Q}_i$ have distinct terminals.

3. For odd-numbered indices, we have the inequality

$$|\mathcal{Q}_{2i+1}| \geq -3k + A\left(\frac{1}{\Delta}\right)^i \prod_{j \leq i}\left(1 - \frac{j}{D}\right). \tag{12}$$

4. For even-numbered indices, we have the inequality

$$e(T(\mathcal{Q}_{2i}), V_2) \geq d|\mathcal{Q}_{2i-1}|. \tag{13}$$

Let

$$t \stackrel{\text{def}}{=} \lceil B/2 \rceil .$$

We will repeatedly use the following straightforward fact, which we call the *small-degree argument*: whenever Q is a good path and $u \in V_2 \setminus Q$ is adjacent to the terminal of Q, then u is adjacent to fewer than t vertices in $V_1 \cap Q$. Indeed, if vertex u were adjacent to $v_{j_1}, v_{j_2}, \ldots, v_{j_t} \in V_1 \cap Q$ with $j_1 < j_2 < \ldots < j_k$, then $v_{j_2} \leftrightsquigarrow u$ (along path Q) and the edge uv_{j_2} would form a cycle of total length at least $2D(t-2)+2 \geq 2D(B/2-2)+2 \stackrel{(10)}{\geq} 2k$. As uv_{j_3} is a chord of the cycle, and u is adjacent to v_{j_1} that is not on the cycle, that would contradict the assumption that G contains no well-placed Θ-graph.

The set $\mathcal{Q}_1$ consists of all paths of the form Pu for $u \in V_2 \setminus P$. Let us check that the preceding conditions hold for $\mathcal{Q}_1$. Vertex v_{l-1} cannot be adjacent to k or more vertices in $P \cap V_2$, for otherwise G would contain a well-placed Θ-graph with a chord through v_{l-1}. So, $|\mathcal{Q}_1| \geq A - k$. Next, consider any $u \in V_2 \setminus P$ that is a neighbor of v_{l-1}. By the small-degree argument vertex u cannot be adjacent to t or more vertices of $P \cap V_1$, and Pu is good.

Suppose $\mathcal{Q}_{2i-1}$ has been defined, and we wish to define $\mathcal{Q}_{2i}$. Consider an arbitrary path $Q = v_0 \leftrightsquigarrow v_1 \leftrightsquigarrow \cdots \leftrightsquigarrow v_{l-1} \leftrightsquigarrow u \in \mathcal{Q}_{2i-1}$. Vertex u cannot have k or more neighbors in $Q \cap V_3$, for otherwise G would contain a well-placed Θ-graph with a chord through u. Hence, there are at least d edges of the form uw, where $w \in V_3 \setminus Q$. As we vary u we obtain a family of at least $d|\mathcal{Q}_{2i-1}|$ paths. We let $\mathcal{Q}_{2i}$ consist of any maximal subfamily of such paths with distinct terminals. The condition (13) follows automatically as each vertex of $T(\mathcal{Q}_{2i-1})$ has at least d neighbors in $T(\mathcal{Q}_{2i})$.

Suppose $\mathcal{Q}_{2i}$ has been defined, and we wish to define $\mathcal{Q}_{2i+1}$. Consider an arbitrary path $Q = v_0 \leftrightsquigarrow v_1 \leftrightsquigarrow \cdots \leftrightsquigarrow v_{l-1} \leftrightsquigarrow u \in \mathcal{Q}_{2i}$. An edge uw is called *long* if $w \in P$, and w is at a distance exceeding $2k$ from u along path Q. If uw is a long edge, then from u to Q there is only one edge, namely the edge to the predecessor of u on Q, for otherwise there is a well-placed Θ-graph. Also, at most i neighbors of u lie on the path $v_{l-1} \leftrightsquigarrow u$. Since $\deg u \geq D$, it follows that there are at least $(1 - i/D) \deg u$ short edges from u that miss $v_{l-1} \leftrightsquigarrow u$. Thus there is a set $\mathcal{W}$ of at least $(1 - i/D)e(T(\mathcal{Q}_{2i}), V_2)$ walks (not necessarily paths!) of the form $v_0 \leftrightsquigarrow v_1 \leftrightsquigarrow \cdots \leftrightsquigarrow v_{l-1} \leftrightsquigarrow uw$ such that $v_{l-1} \leftrightsquigarrow uw$ is a path and w occurs only among the last $2k$ vertices of the walk.

From the maximum degree condition on V_2 it follows that walks in $\mathcal{W}$ have at least $(1 - i/D)e(T(\mathcal{Q}_{2i}), V_2)/\Delta d$ distinct terminals. A walk fails to be a path only if the terminal vertex lies on P. However, since the edge uw is short, this can happen for at most $2k$ possible terminals. Hence, there is a $\mathcal{Q}_{2i+1} \subset \mathcal{W}$ of size

$$|\mathcal{Q}_{2i+1}| \geq (1 - i/D)e(T(\mathcal{Q}_{2i}), V_2)/\Delta d - 2k \tag{14}$$

that consists of paths with distinct terminals. It remains to check that every path in $\mathcal{Q}_{2i+1}$ is good. The only way that $Q = v_0 \leftrightsquigarrow \cdots \leftrightsquigarrow v_{l-1} \leftrightsquigarrow uw \in \mathcal{Q}_{2i+1}$ may fail to be good is if w has no neighbors

in $V_1 \setminus Q$. By the small-degree argument w has fewer than t neighbors in V_1. Since w has at least B neighbors in V_1 and $B \geq t+2$, we conclude that w has at least *two* neighbors in V_1 outside the path. Of course, the same is true for every terminal of a path in $\mathcal{Q}_{2i+1}$. The condition (12) for $\mathcal{Q}_{2i+1}$ follows from (14), (13) and from validity of (12) for $\mathcal{Q}_{2i-1}$.

Note that $\mathcal{Q}_{2D-1}$ is non-empty. Let $Q = v_0 \leftrightsquigarrow \cdots \leftrightsquigarrow v_{l-1} \leftrightsquigarrow u \in \mathcal{Q}_{2D-1}$ be an arbitrary path. Note that since $2D-1$ is odd, $u \in V_2$. By the property of terminals of V_i (odd i) that we noted in the previous paragraph, there are two vertices in $V_1 \setminus Q$ that are neighbors of u. Let v_l be any of them, and let the new path be $Qv_l = v_0 \leftrightsquigarrow \cdots \leftrightsquigarrow v_{l-1} \leftrightsquigarrow uv_l$. This path can fail to be good if there is a vertex w on the path Q that is good in Q, but is bad in Qv_l. By the small-degree argument, w is adjacent to fewer than t vertices in $Q \cap V_1$ that precede w in Q. The same argument applied to the reversal of the path Qv_l shows that w is adjacent to fewer than t vertices in $Q \cap V_1$ that succeeds w in Q. Since $2t-2 < B$, the path Qv_l is good.

Hence, it is possible to build an arbitrarily long path in G. This contradicts the finiteness of G. □

Lemma 6 follows from Lemmas 8 and 9 by setting $C = d+k$, in view of inequality $4k^2+k \leq 5k^2$. We lose k^2-k here for cosmetic reason: $5k^2$ is tidier than $4k^2+k$.

3 Proof of Theorem 1

Suppose that G is a bipartite graph of minimum degree at least $2d+5k^2$ and contains no C_{2k}. Pick a root vertex x arbitrarily, and let $V_0, V_1, \ldots, V_{k-1}$ be the levels obtained from the exploration process in Section 1.

Lemma 10. *For $1 \leq i \leq k-1$, the graph $G[V_{i-1}, V_i, V_{i+1}]$ contains no well-placed Θ-graph.*

Proof. The following proof is almost an exact repetition of the proof of Claim 3.1 from [14] (which is also reproduced as Lemma 11 below).

Suppose, for the sake of contradiction, that a well-placed Θ-graph $F \subset G[V_{i-1}, V_i, V_{i+1}]$ exists. Let $Y = V_i \cap V(F)$. Since F is well-placed, for every vertex of Y there is a path avoiding $V(F)$ of length i to the vertex x. The union of these paths forms a tree T with x as a root. Let y be the vertex farthest from x such that every vertex of Y is a T-descendant of y. Paths that connect x to Y branch at y. Pick one such branch, and let $W \subset Y$ be the set of all the T-descendants of that branch. Let $Z = V(F) \setminus W$. From $W \neq V_i \cap V(F)$ it follows that Z is not an independent set of F, and so $W \cup Z$ is not a bipartition of F.

Let ℓ be the distance between x and y. We have $\ell < i$ and $2k-2i+2\ell < 2k \leq |V(F)|$. By Lemma 3 in F there is a path P of length $2k-2i+2\ell$ that starts at some $w \in W$ and ends in $z \in Z$. Since the length of P is even, $z \in Y$. Let P_w and P_z be unique paths in T that connect y to respectively w and z. They intersect only at y. Each of P_w and P_z has length $i-\ell$. The union of paths P, P_w, P_z forms a $2k$-cycle in G. □

The same argument (with a different Y) also proves the next lemma.

Lemma 11 (Claim 3.1 in [14])**.** *For $1 \leq i \leq k-1$, neither of $G[V_i]$ and $G[V_i, V_{i+1}]$ contains a bipartite Θ-graph.*

The next step is to show that the levels $V_0, V_1, V_2, \ldots$ increase in size. We shall show by induction on i that

$$e(V_i, V_{i+1}) \geq d|V_i|, \tag{15}$$
$$e(V_i, V_{i+1}) \leq 2k|V_{i+1}|, \tag{16}$$
$$e(V_i, V'_{i+1}) \leq 2k|V'_{i+1}|, \tag{17}$$
$$|V_{i+1}| \geq (2k)^{-1} d|V_i|, \tag{18}$$
$$|V_{i+1}| \geq \tfrac{d^2}{400k \log k}|V_{i-1}|. \tag{19}$$

To prove Theorem 1, we only need (19); the remaining inequalities play auxiliary roles in derivation of (19).

Clearly, these inequalities hold for $i = 0$ since each vertex of V_1 sends only one edge to V_0.

Proof of (15): By Lemma 2 the degree of every vertex in V_i is at least $2d + 4k$, and so

$$e(V_i, V'_{i+1}) \geq (2d + 4k)|V_i| - e(V_{i-1}, V_i) \overset{\text{induc.}}{\geq} (2d + 2k)|V_i|.$$

We next distinguish two cases depending on whether V_{i+1} is big (in the sense of the definition from Section 1). If V_{i+1} is big, then $e(V_i, V_{i+1}) = e(V_i, V'_{i+1})$, and (15) follows. If V_{i+1} is normal, then Corollary 5 and Lemma 11 imply that

$$e(V_i, \mathrm{Bg}_{i+1}) \leq k(|V_i| + |\mathrm{Bg}_{i+1}|) \leq k\left(|V_i| + \tfrac{1}{2k}|V'_{i+1}|\right) \leq k|V_i| + \tfrac{1}{2}e(V_i, V'_{i+1})$$

and so

$$e(V_i, V_{i+1}) = e(V_i, V'_{i+1}) - e(V_i, \mathrm{Bg}_{i+1}) \geq \tfrac{1}{2}e(V_i, V'_{i+1}) - k|V_i| \geq d|V_i|$$

implying (15). □

Proof of (16): Consider the graph $G[V_i, V_{i+1}]$. Inequality (15) asserts that the average degree of V_i is at least $d \geq 2k$. If (16) does not hold, then the average degree of V_{i+1} is at least $2k$ as well, contradicting Corollary 5 and Lemma 11. □

Proof for (17): The argument is the same as for (16) with $G[V_i, V'_{i+1}]$ in place of $G[V_i, V_{i+1}]$. □

Proof for (18): This follows from (16) and (15). □

Proof of (19) in the case V_i is a normal level: We assume that (19) does not hold and will derive a contradiction. Consider the trilayered graph $G[V_{i-1}, V_i, V'_{i+1}]$. Let $t = 2 \log k$. Suppose momentarily that the inequalities (2) in Lemma 6 hold. Then since V_i is normal, each vertex in V_i has at most Δd neighbors in V'_{i+1}, and so Lemma 6 applies. However, the lemma's conclusion contradicts Lemmas 10 and 11. Hence, to prove (19) it suffices to verify inequalities (2a–d) with $F = d \cdot e(V_{i-1}, V_i)/8k|V'_{i+1}|$.

We may assume that

$$F \geq 2e^2 \log k, \tag{20}$$

and in particular that (2a) holds. Indeed, if (20) were not true, then inequality (15) would imply $|V'_{i+1}| \geq (d^2/16e^2k\log k)|V_{i-1}|$, and thus

$$|V_{i+1}| \geq (1-\tfrac{1}{k})|V'_{i+1}| \geq (d^2/32e^2k\log k)|V_{i-1}|,$$

and so (19) would follow in view of $32e^2 \leq 400$.

Inequality (2b) is implied by (18). Indeed,

$$e(V_{i-1},V_i) = 8k|V'_{i+1}|F/d \geq 8k|V_{i+1}|F/d \overset{(18)}{\geq} 4F|V_i| \overset{(18)}{\geq} 2k^{-1}dF|V_{i-1}|,$$

and $d \geq k^2$ by the definition of d from (1).

Inequality (2c) is implied by (1) and (15).

Next, suppose (2d) were not true. Since $F/t \geq e^2$ by (20), we would then conclude

$$\begin{aligned}|V_{i+1}| &\overset{(18)}{\geq} (2k)^{-1}d|V_i| \geq d(16k^2)^{-1}(F/et)^t e(V_{i-1},V_i)\\ &\geq d(16k^2)^{-1}e^{2\log k}e(V_{i-1},V_i) \overset{(15)}{\geq} \tfrac{1}{16}d^2|V_{i-1}|,\end{aligned}$$

and so (19) would follow.

Finally, (2e) is a consequence of (15). □

Proof of (19) in the case V_i is a big level: We have

$$\begin{aligned}|V_{i+1}| &\geq \tfrac{1}{2}|V'_{i+1}| \overset{(17)}{\geq} (4k)^{-1}e(V_i,V'_{i+1}) \geq (4k)^{-1}e(\mathrm{Bg}_i,V'_{i+1}) \geq (4k)^{-1}\Delta d|\mathrm{Bg}_i|\\ &\geq (8k^2)^{-1}\Delta d|V_i| \overset{(18)}{\geq} (16k^3)^{-1}\Delta d^2|V_{i-1}| = \tfrac{1}{16}d^2|V_{i-1}|,\end{aligned}$$

and so (19) holds. □

We are ready to complete the proof of Theorem 1. If k is even, then $k/2$ applications of (19) yield

$$|V_k| \geq \frac{d^k}{(400k\log k)^{k/2}}.$$

If k is odd, then $(k-1)/2$ applications of (19) yield

$$|V_k| \geq \frac{d^{k-1}}{(400k\log k)^{(k-1)/2}}|V_1| \geq \frac{d^k}{(400k\log k)^{(k-1)/2}}.$$

Either way, since $|V_k| < n$ we conclude that $d < 20\sqrt{k\log k}\cdot n^{1/k}$.

4 Acknowledgment

We would like to thank the referees for carefully reading the manuscript and for giving constructive comments which helped improving the quality of the paper. All remaining errors are ours.

References

[1] Clark T. Benson. Minimal regular graphs of girths eight and twelve. *Canad. J. Math.*, 18:1091–1094, 1966.

[2] Pavle V. M. Blagojević, Boris Bukh, and Roman Karasev. Turán numbers for $K_{s,t}$-free graphs: topological obstructions and algebraic constructions. *Israel J. Math.*, 197(1):199–214, 2013. arXiv:1108.5254.

[3] J. A. Bondy and M. Simonovits. Cycles of even length in graphs. *J. Combinatorial Theory Ser. B*, 16:97–105, 1974.

[4] W. G. Brown. On graphs that do not contain a Thomsen graph. *Canad. Math. Bull.*, 9:281–285, 1966.

[5] P. Erdős. On sequences of integers no one of which divides the product of two others and on some related problems. *Inst. Math. Mech. Univ. Tomsk*, 2:74–82, 1938.

[6] P. Erdős. Extremal problems in graph theory. In *Theory of Graphs and its Applications (Proc. Sympos. Smolenice, 1963)*, pages 29–36. Publ. House Czechoslovak Acad. Sci., Prague, 1964.

[7] P. Erdős and A. Rényi. On a problem in the theory of graphs. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 7:623–641 (1963), 1962.

[8] Zoltan Füredi, Assaf Naor, and Jacques Verstraëte. On the Turán number for the hexagon. *Adv. Math.*, 203(2):476–496, 2006. https://web.math.princeton.edu/~naor/homepagefiles/final-hexagons.pdf.

[9] Zoltán Füredi and Miklós Simonovits. The history of degenerate (bipartite) extremal graph problems. arXiv:1306.5167, June 2013.

[10] Peter Keevash. Hypergraph Turán problems. In *Surveys in combinatorics 2011*, volume 392 of *London Math. Soc. Lecture Note Ser.*, pages 83–139. Cambridge Univ. Press, Cambridge, 2011. http://people.maths.ox.ac.uk/keevash/papers/turan-survey.pdf.

[11] T. Kövari, V. T. Sós, and P. Turán. On a problem of K. Zarankiewicz. *Colloquium Math.*, 3:50–57, 1954.

[12] Felix Lazebnik and Vasiliy A. Ustimenko. Explicit construction of graphs with an arbitrary large girth and of large size. *Discrete Appl. Math.*, 60(1-3):275–284, 1995. ARIDAM VI and VII (New Brunswick, NJ, 1991/1992).

[13] Keith E. Mellinger and Dhruv Mubayi. Constructions of bipartite graphs from finite geometries. *J. Graph Theory*, 49(1):1–10, 2005. http://homepages.math.uic.edu/~mubayi/papers/ArcConstf.pdf.

[14] Oleg Pikhurko. A note on the Turán function of even cycles. *Proc. Amer. Math. Soc.*, 140(11):3687–3692, 2012. http://homepages.warwick.ac.uk/~maskat/Papers/EvenCycle.pdf.

[15] Alexander Sidorenko. What we know and what we do not know about Turán numbers. *Graphs Combin.*, 11(2):179–199, 1995.

[16] M. Simonovits. A method for solving extremal problems in graph theory, stability problems. In *Theory of Graphs (Proc. Colloq., Tihany, 1966)*, pages 279–319. Academic Press, New York, 1968.

[17] P. Turán. On an extremal problem in graph theory (in Hungarian). *Mat. Fiz. Lapok*, 48:436–452, 1941.

[18] Jacques Verstraëte. On arithmetic progressions of cycle lengths in graphs. *Combin. Probab. Comput.*, 9(4):369–373, 2000. arXiv:math/0204222.

[19] R. Wenger. Extremal graphs with no C^4's, C^6's, or C^{10}'s. *J. Combin. Theory Ser. B*, 52(1):113–116, 1991.